

Table 4b. Site-wise number of cases and relative survival by age group: Cuba, 1994–1995 cases followed-up until 1999

Site	ICD-10	Cases included	Number of cases by age group					Relative survival by age group				
								% 5-year survival				
			< 45	45-54	55-64	65-74	> 75	< 45	45-54	55-64	65-74	> 75
Tongue	C01-02	231	18	30	59	60	64	50.5	37.5	34.9	37.2	32.8
Oral cavity	C03-06	260	12	35	67	72	74	51.0	41.2	47.3	36.6	39.8
Tonsil	C09	54	8	11	8	18	9	76.0	37.7	40.3	64.9	0.0
Oropharynx	C10	48	1	9	15	14	9	100.8	68.7	21.4	55.2	33.6
Colon	C18	1 248	74	124	247	345	458	51.8	40.8	43.3	49.7	31.7
Rectum	C19-20	556	40	79	113	167	157	52.0	52.9	52.6	47.5	42.9
Anus	C21	94	5	17	22	20	30	60.6	46.0	46.6	84.0	58.8
Larynx	C32	800	50	130	238	244	138	80.7	63.5	49.5	51.3	38.9
Breast	C50	2 169	424	515	502	435	293	71.7	67.0	70.7	72.6	65.4
Cervix	C53	1 221	531	284	184	155	67	65.3	55.3	46.7	54.5	53.0
Urinary bladder	C67	819	31	82	157	241	308	74.4	80.6	71.4	59.4	57.2
Hodgkin lymphoma	C81	186	108	15	22	25	16	63.6	61.4	43.9	18.4	20.3
Non-Hodgkin lymphoma	C82-85+C96	464	122	80	87	97	78	56.7	54.9	41.8	37.5	43.4