

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Barshi, India, 1993–2000 cases followed-up until 2003

Site	ICD-10	Cases included	Complete FU		Incomplete FU: lost to FU						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Tongue	C01-02	47	47	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	9.5
Oral cavity	C03-06	55	53	96.4	2	3.6	0.0	1.8	1.8	0.0	96.4	10.1
Hypopharynx	C12-13	80	79	98.8	1	1.3	0.0	0.0	1.3	0.0	98.8	7.5
Oesophagus	C15	97	96	99.0	1	1.0	0.0	0.0	0.0	1.0	100.0	3.9
Stomach	C16	45	43	95.6	2	4.4	2.2	0.0	2.2	0.0	95.6	2.8
Rectum	C19-20	49	48	98.0	1	2.0	2.0	0.0	0.0	0.0	98.0	9.4
Liver	C22	47	47	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	1.6
Larynx	C32	33	32	97.0	1	3.0	3.0	0.0	0.0	0.0	97.0	12.1
Lung	C33-34	47	45	95.7	2	4.3	2.1	2.1	0.0	0.0	95.7	3.4
Other skin	C44	39	39	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	49.2
Breast	C50	124	119	96.0	5	4.0	0.0	0.0	3.2	0.8	96.8	41.4
Cervix	C53	406	403	99.3	3	0.7	0.2	0.0	0.2	0.2	99.5	24.4
Penis	C60	43	42	97.7	1	2.3	0.0	0.0	2.3	0.0	97.7	38.3
Non-Hodgkin lymphoma	C82-85+C96	36	36	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	9.8
Myeloid leukaemia	C92-94	40	40	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	4.0

FU: follow-up