

Table 3. Comparison of 1-, 3- and 5-year absolute and relative survival and 5-year age-standardized relative survival by site: Bhopal, India, 1991–1995 cases followed-up until 2000

Site	ICD-10	Cases included	% Absolute survival			% Relative survival			% ASRS at 5-years	
			1-year	3-year	5-year	1-year	3-year	5-year	all ages	0-74 years
Tongue	C01-02	127	52.8	20.5	9.4	54.8	22.6	10.8	11.7	13.1
Oral cavity	C03-06	172	80.2	45.9	29.1	83.0	50.4	33.6	33.9	37.5
Hypopharynx	C12-13	118	50.8	16.9	1.7	52.7	18.6	2.0	1.7	2.0
Oesophagus	C15	163	36.2	12.9	3.1	37.6	14.1	3.5	2.7	3.6
Stomach	C16	75	38.7	5.3	2.7	39.8	5.8	2.9	2.8	3.8
Colon	C18	48	52.1	25.0	6.3	53.9	27.1	7.0	3.2	5.2
Rectum	C19-20	39	56.4	25.6	7.7	58.0	27.2	8.5	4.0	6.5
Larynx	C32	60	61.7	23.3	6.7	63.6	25.4	7.8	12.5	15.7
Lung	C33-34	216	33.3	6.9	0.9	34.7	7.7	1.1	0.8	1.0
Breast	C50	261	83.1	62.1	30.7	84.5	64.8	32.8	25.3	30.6
Cervix	C53	332	88.9	66.0	33.1	90.3	68.8	35.4	30.8	34.5
Ovary	C56	69	63.8	47.8	18.8	64.8	49.7	20.1	15.8	18.9
Urinary bladder	C67	46	54.3	17.4	6.5	57.1	19.6	7.8	6.0	9.7
Non-Hodgkin lymphoma	C82-85+C96	30	33.3	16.7	10.0	33.9	17.3	10.5	8.8	9.9
Lymphoid leukaemia	C91	32	46.9	31.3	12.5	47.5	32.1	13.1	10.4	12.6
Myeloid leukaemia	C92-94	75	61.3	33.3	16.0	62.0	34.3	16.7	12.3	14.2

ASRS: age-standardized relative survival