

**Table 4a. Site-wise number of cases, 5-year absolute and relative survival by sex: Chennai, India, 1990–1999 cases followed-up to 2001**

Site	ICD-10	Cases included	Male			Female		
			% 5-year survival			% 5-year survival		
			No.	Abs	Rel	No.	Abs	Rel
Lip	C00	86	47	46.3	52.4	39	34.4	41.3
Tongue	C01-02	988	761	18.1	21.8	227	23.9	26.6
Oral cavity	C03-06	1 662	874	30.9	37.0	788	30.1	34.2
Tonsil	C09	250	213	11.2	14.0	37	27.2	31.6
Oropharynx	C10	158	135	15.9	19.1	23	13.0	15.4
Hypopharynx	C12-13	1 017	757	11.9	14.2	260	14.2	15.6
Oesophagus	C15	2 016	1 229	5.6	7.0	787	9.2	10.5
Stomach	C16	2 681	1 828	8.3	10.0	853	9.2	10.3
Pancreas	C25	328	231	6.8	7.7	97	10.7	12.2
Larynx	C32	722	651	30.8	37.1	71	30.0	33.6
Lung	C33-34	1 806	1 527	6.1	7.2	279	8.7	9.7
Breast	C50	3 067				3 067	43.7	48.6
Cervix	C53	4 438				4 438	54.0	59.4
Ovary	C56	808				808	27.4	29.7
Urinary bladder	C67	442	318	23.2	30.6	124	23.6	27.9
Hodgkin lymphoma	C81	298	213	36.8	38.5	85	45.5	46.9
Non-Hodgkin lymphoma	C82-85+C96	868	558	20.1	22.1	310	24.5	26.3
Lymphoid leukaemia	C91	433	282	23.1	23.9	151	25.3	25.9
Myeloid leukaemia	C92-94	465	262	14.2	15.4	203	15.3	16.2
Leukaemia unspecified	C95	85	50	10.5	11.1	35	11.2	11.9

*Abs: absolute survival; Rel: relative survival*