

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Khon Kaen, Thailand, 1993–1997 cases followed-up until 2000

Site	ICD-10	No. of cases included	Complete FU		Incomplete FU: loss to FU*						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Lip	C00	79	53	67.1	26	32.9	19.0	5.1	6.3	2.5	69.6	32.0
Tongue	C01-02	54	42	77.8	12	22.2	20.4	1.9	0.0	0.0	77.8	8.4
Oral cavity	C03-06	107	79	73.8	28	26.2	19.6	2.8	1.9	1.9	75.7	12.3
Nasopharynx	C11	118	95	80.5	23	19.5	12.7	4.2	0.8	1.7	82.2	18.1
Colon	C18	242	193	79.8	49	20.2	6.6	6.2	5.4	2.0	81.8	18.2
Rectum	C19-20	139	112	80.6	27	19.4	10.1	6.5	0.7	2.1	82.7	18.6
Larynx	C32	36	29	80.6	7	19.4	19.4	0.0	0.0	0.0	80.6	13.8
Breast	C50	418	173	41.4	245	58.6	24.4	22.2	7.7	4.3	45.7	21.5
Cervix	C53	554	210	37.9	344	62.1	39.2	15.9	5.1	1.9	39.8	9.9
Ovary	C56	206	97	47.1	109	52.9	28.6	12.6	8.7	3.0	50.1	11.5
Urinary bladder	C67	103	43	41.7	60	58.3	35.9	10.7	8.7	3.0	44.7	7.8
Hodgkin lymphoma	C81	29	11	37.9	18	62.1	27.6	17.2	6.9	10.4	48.3	24.3
Non-Hodgkin lymphoma	C82-85+C96	168	89	53.0	79	47.0	28.0	10.1	6.0	2.9	55.9	9.1

FU: follow-up; * non-random