

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Izmir, Turkey, 1995–1997 cases followed-up until 2003

Site	ICD-10	Cases included	Complete FU		Incomplete FU: lost to FU						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Colon	C18	371	266	71.7	105	28.3	11.1	3.0	1.3	12.9	84.6	28.2
Rectum	C19-20	344	232	67.5	112	32.5	16.0	2.9	2.0	11.6	79.1	24.6
Larynx	C32	478	308	64.4	170	35.6	9.8	1.7	1.5	22.6	87.0	64.0
Breast	C50	1 329	806	60.7	523	39.3	10.9	2.3	1.8	24.3	85.0	72.0
Cervix	C53	246	127	51.6	119	48.4	10.6	3.7	2.4	31.7	83.3	51.9
Ovary	C56	230	141	61.3	89	38.7	10.9	1.3	2.6	23.9	85.2	49.6
Urinary bladder	C67	566	350	61.8	216	38.2	13.3	2.7	1.2	21.0	82.9	49.3
Hodgkin lymphoma	C81	85	56	65.8	29	34.2	11.8	5.9	1.2	15.3	81.1	65.0
Non-Hodgkin lymphoma	C82-85+C96	344	256	74.5	88	25.5	11.6	3.2	2.0	8.7	83.2	22.4
Multiple myeloma	C90	79	68	86.0	11	14.0	7.6	1.3	1.3	3.8	89.8	19.4
Lymphoid leukaemia	C91	161	99	61.4	62	38.6	7.5	2.5	1.9	26.7	97.8	31.9
Myeloid leukaemia	C92-94	148	119	80.3	29	19.7	6.8	4.1	2.0	6.8	87.1	16.9

FU: follow-up