

Table 3. Comparison of 1-, 3- and 5-year absolute and relative survival and 5-year age-standardized relative survival by site: Izmir, Turkey, 1995–1997 cases followed-up until 2003

Site	ICD-10	Cases included	% Absolute survival			% Relative survival			% ASRS at 5-years	
			1-year	3-year	5-year	1-year	3-year	5-year	all ages	0-74 years
Colon	C18	371	73.8	53.6	44.9	76.0	58.7	52.5	53.2	52.2
Rectum	C19-20	344	82.9	55.9	44.1	85.0	60.1	50.4	51.6	52.0
Larynx	C32	478	87.9	70.2	62.2	90.0	75.6	70.9	68.9	71.4
Breast	C50	1 329	96.3	81.3	71.1	97.6	84.9	76.7	76.6	77.2
Cervix	C53	246	89.7	68.2	58.8	90.7	70.5	62.3	57.6	63.5
Ovary	C56	230	83.4	62.9	54.7	84.7	65.7	59.0	60.2	59.7
Urinary bladder	C67	566	84.7	67.3	57.7	87.7	75.1	69.7	69.9	70.7
Hodgkin lymphoma	C81	85	92.5	77.4	65.6	93.3	79.3	68.6	67.9	65.8
Non-Hodgkin lymphoma	C82-85+C96	344	72.6	51.0	45.1	73.9	53.8	49.5	49.8	50.6
Multiple myeloma	C90	79	65.8	43.0	28.5	67.3	46.2	32.3	31.6	31.1
Lymphoid leukaemia	C91	161	78.1	55.5	48.2	79.2	57.6	51.4	52.6	50.1
Myeloid leukaemia	C92-94	148	65.7	41.2	29.5	67.0	43.4	32.2	33.0	34.2

ASRS: age-standardized relative survival