

Table 4a. Site-wise number of cases, 5-year absolute and relative survival by sex: Izmir, Turkey, 1995–1997 cases followed-up until 2003

Site	ICD-10	Cases included	Male			Female		
			% 5-year survival			% 5-year survival		
			No.	Abs	Rel	No.	Abs	Rel
Colon	C18	371	217	42.8	50.1	154	48.0	56.1
Rectum	C19-20	344	192	47.8	55.9	152	39.6	43.8
Larynx	C32	478	450	61.9	70.8	28	66.2	73.2
Breast	C50	1 329	36	66.5	76.4	1 293	71.2	76.7
Cervix	C53	246				246	58.8	62.3
Ovary	C56	230				230	54.7	59.0
Urinary bladder	C67	566	508	57.5	69.4	58	59.2	72.0
Hodgkin lymphoma	C81	85	56	65.1	68.4	29	66.4	68.8
Non-Hodgkin lymphoma	C82-85+C96	344	194	40.3	43.9	150	51.4	56.7
Multiple myeloma	C90	79	41	26.0	30.4	38	31.5	34.6
Lymphoid leukaemia	C91	161	93	45.8	49.6	68	51.3	53.8
Myeloid leukaemia	C92-94	148	78	23.4	26.2	70	38.0	40.4

*Abs: absolute survival; Rel: relative survival*