

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Kampala, Uganda, 1993–1997 cases followed-up until 1999

Site	ICD-10	No. of cases included	Complete FU		Incomplete FU: loss to FU*						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Nasopharynx	C11	49	31	63.3	18	36.7	32.7	4.1	0.0	0.0	63.3	6.3
Oesophagus	C15	180	152	84.4	28	15.6	12.8	2.8	0.0	0.0	84.4	7.2
Stomach	C16	90	67	74.4	23	25.6	21.1	4.4	0.0	0.0	74.4	4.2
Colon	C18	51	41	80.4	10	19.6	19.6	0.0	0.0	0.0	80.4	6.6
Rectum	C19-20	50	40	80.0	10	20.0	18.0	2.0	0.0	0.0	80.0	9.4
Liver	C22	116	101	87.1	15	12.9	12.1	0.0	0.9	0.0	87.1	5.9
Lung	C33-34	47	41	87.2	6	12.8	12.8	0.0	0.0	0.0	87.2	4.8
Kaposi sarcoma	C46 ^o	370	248	67.0	122	33.0	21.6	9.2	1.9	0.3	67.3	8.1
Breast	C50	162	96	59.3	66	40.7	27.8	9.3	1.9	1.9	61.1	10.9
Cervix	C53	283	149	52.7	134	47.3	35.0	11.3	0.7	0.4	53.0	8.3
Ovary	C56	66	48	72.7	18	27.3	15.2	7.6	3.0	1.5	74.2	5.0
Prostate	C61	154	119	77.3	35	22.7	16.2	3.9	2.6	0.0	77.3	26.2
Eye	C69	81	38	46.9	43	53.1	37.0	14.8	1.2	0.0	46.9	14.9
Thyroid	C73	39	22	56.4	17	43.6	25.6	15.4	0.0	2.6	59.0	7.5
Non-Hodgkin lymphoma	C82-85+C96	178	118	66.3	60	33.7	23.0	7.9	1.7	1.1	67.4	9.4

FU: follow-up; ^o from a random sample of total cases; * non-random