

Table 3. Comparison of 1-, 3- and 5-year absolute and relative survival and 5-year age-standardized relative survival by site: Kampala, Uganda, 1993–1997 cases followed-up until 1999

Site	ICD-10	Cases included	% Absolute survival			% Relative survival			% ASRS at 5-years	
			1-year	3-year	5-year	1-year	3-year	5-year	all ages	0-74 years
Nasopharynx	C11	49	58.5	27.2	0.0	59.8	29.0	0.0	0.0	0.0
Oesophagus	C15	180	36.5	10.3	2.5	38.3	12.3	3.4	4.6	3.0
Stomach	C16	90	39.1	7.5	0.0	40.9	8.5	0.0	0.0	0.0
Colon	C18	51	54.3	19.0	5.4	57.2	22.7	7.4	6.2	8.2
Rectum	C19-20	50	56.0	13.7	9.1	58.4	15.4	11.0	7.2	10.1
Liver	C22	116	33.0	8.0	2.7	34.0	8.7	3.1	1.1	1.4
Lung	C33-34	47	15.9	0.0	0.0	16.4	0.0	0.0	0.0	0.0
Kaposi sarcoma	C46*	370	58.5	33.8	18.6	59.7	36.1	20.8	21.9	35.6
Breast	C50	162	72.0	54.3	37.8	74.5	59.3	44.0	36.1	45.8
Cervix	C53	283	73.0	34.9	17.1	75.0	37.8	19.8	18.7	13.1
Ovary	C56	66	44.3	28.5	13.7	45.5	31.4	16.1	10.2	8.6
Prostate	C61	154	86.6	56.2	29.4	95.0	75.3	48.1	45.6	47.2
Eye	C69	81	92.4	55.1	30.4	94.3	58.4	33.4	35.7	35.7
Thyroid	C73	39	62.3	26.7	8.9	63.8	28.7	10.2	10.8	10.8
Non-Hodgkin lymphoma	C82-85+C96	178	61.4	40.7	31.9	62.3	42.2	33.6	26.1	25.1

ASRS: age-standardized relative survival; * random sample of total incident cases