

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Harare, Zimbabwe, 1993–1997 cases* followed-up until 1999, all races together

Site	ICD-10	No. of cases included	Complete FU		Incomplete FU: loss to FU*						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Oesophagus	C15	131	131	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	2.3
Stomach	C16	129	127	98.4	2	1.6	1.6	0.0	0.0	0.0	98.4	4.8
Colon	C18	57	57	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	14.6
Rectum	C19-20	67	65	97.0	2	3.0	1.5	1.5	0.0	0.0	97.0	16.1
Liver	C22	132	131	99.2	1	0.8	0.8	0.0	0.0	0.0	99.2	1.0
Larynx	C32	73	69	94.5	4	5.5	5.5	0.0	0.0	0.0	94.5	14.1
Lung	C33-34	116	116	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	2.0
Melanoma of skin	C43	101	100	99.0	1	1.0	1.0	0.0	0.0	0.0	99.0	54.2
Kaposi sarcoma	C46	151	151	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	3.4
Breast	C50	258	250	96.9	8	3.1	2.7	0.4	0.0	0.0	96.9	39.3
Cervix	C53	264	260	98.5	4	1.5	1.1	0.4	0.0	0.0	98.5	26.2
Ovary	C56	63	63	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	18.7
Prostate	C61	64	61	95.3	3	4.7	4.7	0.0	0.0	0.0	95.3	16.6
Urinary bladder	C67	130	127	97.7	3	2.3	1.5	0.8	0.0	0.0	97.7	11.8
Eye	C69	116	114	98.3	2	1.7	0.9	0.9	0.0	0.0	98.3	42.1
Hodgkin lymphoma	C81	39	39	100.0	0	0.0	0.0	0.0	0.0	0.0	100.0	30.6
Non-Hodgkin lymphoma	C82-85+C96	99	97	98.0	2	2.0	2.0	0.0	0.0	0.0	98.0	7.8

FU: follow-up; * from a random sample of total cases