

Table 4c. Frequency and 5-year absolute survival by clinical extent of disease for all registries: cancer of the larynx

Country/ registry	Period of registration	Frequency (%)				5-year absolute survival (%)			
		L	R	D	U	L	R	D	U
Cuba	1994–1995	51.3	23.0	1.3	24.5	65.5	34.5	10.0	22.2
India									
Chennai	1990–1999	6.6	85.5	3.6	4.3	44.3	30.3	-	34.1
Karunagappally	1991–1997	30.4	45.7	13.0	10.9	55.6	19.0	0.0	-
Mumbai	1992–1999	33.2	51.1	9.7	6.1	57.1	16.5	0.5	23.2
Singapore	1993–1997	35.7	12.2	2.3	49.8	63.6	28.6	14.3	41.8
Thailand									
Chiang Mai	1993–1997	11.8	82.4	3.8	2.0	52.4	21.4	16.7	0.0
Lampang	1990–2000	17.3	61.7	7.4	13.6	53.4	31.6	25.0	45.5
Songkhla	1990–1999	21.0	39.9	8.7	30.4	59.5	31.9	42.4	21.9
Turkey Izmir	1995–1997	32.1	23.0	7.5	37.4	74.8	45.1	47.6	65.8

L: localized; R: regional; D: distant metastasis; U: unknown