

Table 4g. Frequency and 5-year absolute survival by clinical extent of disease for all registries: cancer of the cervix

Country/ registry	Period of registration	Frequency (%)				5-year absolute survival (%)			
		L	R	D	U	L	R	D	U
Costa Rica	1995–2000	22.4	40.5	4.0	33.1	89.5	43.1	11.3	43.2
Cuba	1994–1995	41.3	34.3	1.7	22.7	73.9	41.5	33.3	45.0
India									
Bhopal	1991–1995	28.3	70.5	0.3	0.9	60.6	22.7	0.0	0.0
Chennai	1990–1999	6.4	86.0	3.7	3.9	69.1	55.3	12.4	43.4
Karunagappally	1991–1997	15.3	60.6	8.8	15.3	72.1	43.5	23.1	44.3
Mumbai	1992–1999	27.9	56.8	8.6	6.7	68.3	35.7	3.4	40.7
Philippines									
Manila	1994–1995	21.5	30.5	10.3	37.7	63.1	29.9	7.1	28.2
Singapore	1993–1997	45.5	5.7	5.0	43.8	69.7	48.0	20.4	55.7
Thailand									
Chiang Mai	1993–1997	26.1	69.7	3.7	0.5	81.2	52.7	12.2	75.0
Khon Kaen	1993–1997	17.3	53.8	6.3	22.6	65.1	48.7	30.6	57.0
Lampang	1990–2000	31.2	53.9	5.8	9.2	78.7	57.9	6.5	70.6
Songkhla	1990–1999	22.3	54.6	5.8	17.3	81.2	56.3	15.4	61.3
Turkey									
Izmir	1995–1997	41.8	41.8	6.1	23.2	67.7	54.6	9.3	69.1

L: localized; R: regional; D: distant metastasis; U: unknown