

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Cuba, 1994–1995 cases followed-up until 1999

Site	ICD-10	Cases included	Complete FU		Incomplete FU: lost to FU						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Tongue	C01-02	231	220	95.2	11	4.8	3.0	0.9	0.9	0.0	95.2	13.2
Oral cavity	C03-06	260	243	93.4	17	6.6	3.1	1.9	1.2	0.4	93.8	19.2
Tonsil	C09	54	53	98.1	1	1.9	0.0	0.0	1.9	0.0	98.1	20.5
Oropharynx	C10	48	47	97.9	1	2.1	2.1	0.0	0.0	0.0	97.9	13.8
Colon	C18	1 248	1 238	99.2	10	0.8	0.2	0.2	0.3	0.1	99.3	14.4
Rectum	C19-20	556	545	98.0	11	2.0	1.1	0.4	0.5	0.0	98.0	28.9
Anus	C21	94	93	98.9	1	1.1	1.1	0.0	0.0	0.0	98.9	50.7
Larynx	C32	800	771	96.3	29	3.7	1.5	0.9	1.0	0.3	96.6	40.6
Breast	C50	2 169	2 104	97.0	65	3.0	0.9	0.3	1.4	0.4	97.4	54.5
Cervix	C53	1 221	1 152	94.4	69	5.6	2.2	1.2	1.7	0.5	94.9	50.4
Urinary bladder	C67	819	798	97.4	21	2.6	1.2	0.0	1.0	0.4	97.8	50.1
Hodgkin lymphoma	C81	186	181	97.4	5	2.6	1.6	0.5	0.5	0.0	97.4	49.2
Non-Hodgkin lymphoma	C82-85+C96	464	453	97.7	11	2.3	0.6	0.6	1.1	0.0	97.7	25.1

FU: follow-up

