

Table 2. Number and proportion of cases with complete/incomplete follow-up (in years) and median follow-up (in months) by site: Chennai, India, 1990–1999 cases followed-up to 2001

Site	ICD-10	Cases included	Complete FU		Incomplete FU: lost to FU						% with complete FU at 5 years	Median FU (in months)
			Alive/dead at end of FU		% lost to FU: years from diagnosis							
			No.	%	No.	%	< 1	1-3	3-5	> 5		
Lip	C00	86	56	65.1	30	34.9	7.0	2.3	0.0	25.6	90.7	27.3
Tongue	C01-02	988	747	75.6	241	24.4	13.1	2.6	1.2	7.5	83.1	9.8
Oral cavity	C03-06	1 662	1 221	73.5	441	26.5	10.3	2.2	1.8	12.2	85.6	12.9
Tonsil	C09	250	230	92.0	20	8.0	4.8	0.8	0.0	2.4	94.4	10.3
Oropharynx	C10	158	151	95.6	7	4.4	0.6	0.6	0.0	3.2	98.7	11.0
Hypopharynx	C12-13	1 017	892	87.7	125	12.3	9.0	0.7	0.3	2.3	90.0	8.7
Oesophagus	C15	2 016	1 818	90.2	198	9.8	6.7	0.9	0.4	1.8	92.0	6.9
Stomach	C16	2 681	2 397	89.4	284	10.6	7.3	0.9	0.9	1.5	90.9	6.1
Pancreas	C25	328	314	95.7	14	4.3	3.1	0.3	0.3	0.6	96.3	4.3
Larynx	C32	722	598	82.8	124	17.2	6.7	0.8	0.3	9.4	92.2	17.4
Lung	C33-34	1 806	1 619	89.6	187	10.4	8.0	0.7	0.3	1.4	91.1	5.8
Breast	C50	3 067	2 351	76.7	716	23.3	12.4	2.9	2.0	6.0	82.6	26.9
Cervix	C53	4 438	2 752	62.0	1 686	38.0	11.0	3.7	2.5	20.8	82.8	27.5
Ovary	C56	808	625	77.4	183	22.6	14.7	4.6	1.4	1.9	79.2	11.4
Urinary bladder	C67	442	367	83.0	75	17.0	10.9	1.6	0.2	4.3	87.3	11.7
Hodgkin lymphoma	C81	298	245	82.2	53	17.8	6.4	1.7	1.0	8.7	90.9	21.0
Non-Hodgkin lymphoma	C82-85+C96	868	732	84.3	136	15.7	10.9	1.3	0.6	2.9	87.2	9.4
Lymphoid leukaemia	C91	433	372	85.9	61	14.1	2.8	3.2	3.5	4.6	90.5	8.5
Myeloid leukaemia	C92-94	465	400	86.0	65	14.0	8.6	1.3	0.4	3.7	89.7	4.5
Leukaemia unspecified	C95	85	74	87.1	11	12.9	10.5	0.0	0.0	2.4	89.4	1.5

FU: follow-up