

**Table 4a. Site-wise number of cases, 5-year absolute and relative survival by sex: Mumbai, India, 1992–1994 cases followed through 1999 and 1995–1999 cases followed-up until 2003**

Site	ICD-10	Cases included	Male			Female		
			% 5-year survival			% 5-year survival		
			No.	Abs	Rel	No.	Abs	Rel
Lip	C00	161	96	52.8	59.7	65	60.4	69.7
Tongue	C01-02	2 106	1 550	22.5	26.2	556	33.0	37.6
Oral cavity	C03-06	2 769	1 768	33.9	38.9	1 001	29.5	33.7
Tonsil	C09	513	453	11.7	14.0	60	30.8	37.9
Oropharynx	C10	243	220	15.3	18.0	23	19.1	24.3
Nasopharynx	C11	243	180	21.7	24.3	63	26.2	28.7
Hypopharynx	C12-13	1 902	1 522	17.1	21.2	380	20.7	23.7
Oesophagus	C15	3 128	1 866	13.5	16.3	1 262	12.3	14.1
Stomach	C16	2 205	1 507	13.1	15.4	698	12.1	13.7
Colon	C18	1 578	918	28.5	34.1	660	26.0	29.8
Rectum	C19-20	1 217	759	28.9	34.2	458	28.2	32.7
Anus	C21	220	127	32.3	37.9	93	36.3	41.6
Pancreas	C25	917	578	12.5	14.6	339	12.5	14.7
Larynx	C32	1 880	1 630	28.4	34.6	250	29.9	34.2
Lung	C33-34	3 451	2 705	11.0	13.5	746	10.5	12.0
Breast	C50	7 294				7 294	46.0	51.4
Cervix	C53	4 436				4 436	42.2	46.1
Ovary	C56	2 029				2 029	22.8	24.6
Penis	C60	327	327	43.6	51.6			
Prostate	C61	1 463	1 463	24.0	35.9			
Testis	C62	383	383	53.0	55.2			
Kidney	C64	819	569	30.1	35.3	250	33.7	37.4
Urinary bladder	C67	1 307	1 046	32.0	41.4	261	34.6	43.3
Hodgkin lymphoma	C81	528	370	53.3	56.5	158	38.9	41.1
Non-Hodgkin lymphoma	C82-85+C96	2 338	1 521	31.7	35.8	817	27.5	30.9
Lymphoid leukaemia	C91	1 099	725	14.6	15.5	374	16.4	17.7
Myeloid leukaemia	C92-94	1 261	755	14.7	15.8	506	12.2	13.0
Leukaemia unspecified	C95	345	195	5.8	6.3	150	6.6	7.2

*Abs: absolute survival; Rel: relative survival*