

Table 4b. Site-wise number of cases and relative survival by age group: Izmir, Turkey, 1995–1997 cases followed-up until 2003

Site	ICD-10	Cases included	Number of cases by age group					Relative survival by age group				
								% 5-year survival				
			< 45	45-54	55-64	65-74	> 75	< 45	45-54	55-64	65-74	> 75
Colon	C18	371	49	60	97	120	45	48.0	57.8	51.8	51.4	55.6
Rectum	C19-20	344	64	66	85	99	30	39.4	48.3	47.7	64.5	53.0
Larynx	C32	478	55	118	164	121	20	72.7	68.9	72.9	71.3	58.4
Breast	C50	1 329	387	360	310	211	61	73.1	75.2	80.4	82.5	72.2
Cervix	C53	246	66	71	66	39	4	70.9	63.3	60.0	54.4	0.0
Ovary	C56	230	58	47	66	48	11	66.9	64.2	49.5	57.1	64.6
Urinary bladder	C67	566	35	85	168	197	81	84.7	73.3	66.6	68.1	67.0
Hodgkin lymphoma	C81	85	58	11	9	5	2	72.5	53.7	62.8	32.7	138.2
Non-Hodgkin lymphoma	C82-85+C96	344	117	55	82	72	18	54.1	50.5	49.4	41.0	42.2
Multiple myeloma	C90	79	9	22	25	20	3	14.0	39.7	34.4	27.3	54.3
Lymphoid leukaemia	C91	161	96	20	16	19	10	53.4	47.1	49.8	37.7	73.7
Myeloid leukaemia	C92-94	148	60	20	22	32	14	38.7	37.4	22.3	23.2	25.3